

Medication Information

Acetaminophen with Oxycodone

(A – seat – a min – oh – fen
with oks – I – KOE – done)

Other names for this medication

Endocet[®]
Oxycocet[®]
Percocet[®]

How this medication is used

This medication is used to treat moderate to moderately severe pain.

How to take this medication

Take this medication exactly as directed by your doctor or health care provider. Usually 1 to 2 tablets can be taken every 4 to 6 hours. Never take more than 4000 mg of acetaminophen a day.

Many other pain, cold and flu medications also contain acetaminophen. Talk to your pharmacist first to make sure you are not taking too much.

This medication does not mix well with alcohol in your body. Ask your doctor or health care provider about drinking alcohol such as wine, beer and liquor while taking this medication.

Oxycodone makes some people drowsy. Do not drive or operate machinery until you know how this medication affects you.

While taking this medication you may notice

- drowsiness
- constipation
- dizziness
- nausea and vomiting
- confusion
- restlessness
- trouble sleeping

To avoid feeling dizzy, get up slowly when you have been sitting or lying down. If you feel dizzy, sit or lie down until you are no longer dizzy, then get up slowly.

To avoid constipation, eat food high in fibre, drink 6 to 8 glasses of liquid each day, and exercise regularly. Food high in fibre includes whole grain products, vegetables and fruit.

If you have stomach upset, take this medication with food. Your doctor or health care provider may suggest another medication to help with any constipation and nausea.

Contact your doctor or health care provider if you notice

- your pain does not get better
- breathing problems
- fainting
- unusual bruising
- severe constipation
- you cannot pass urine
- you are very nervous or excited
- black tarry stools
- continued nausea and vomiting
- yellow eyes or skin